

Linzer biol. Beitr.	23/2	615-620	30.12.1991
---------------------	------	---------	------------

***HELIOTROPIMUM ESFANDIARII* (BORAGINACEAE), A NEW SPECIES**

**FROM NORTH-CENTRAL IRAN**

H. AKHANI, Tehran & H. RIEDL, Wien

**A b s t r a c t :** *Heliotropium esfandiarii* AKHANI & RIEDL is described as a new species from north-central Iran. The new species is related to *H. szovitsii*. Its illustration and distribution map is provided. Recently the first author who is revising the genus *Heliotropium* in Iran and adjacent parts came across some specimens from north-central Iran (northwestern border of Dashte-Kavir), which seem to be different from the other species. Hitherto two specimens of the relevant plants have been seen by the second author. These were identified as *H. szovitsii* (STEV.) BUNGE (RIEDL 1967). Further experiences in field and study of type and several specimens of *H. szovitsii* revealed that the material in question is not only different by taxonomic characters from *H. szovitsii*, but also in geographical distribution. After a joint discussion in Wien we decided to describe these specimens as a new species *H. esfandiarii*, in honour of the Iranian botanist Dr. E. Esfandiari.

***Heliotropium esfandiarii* AKHANI & H. RIEDL spec.nov. (Fig.1)**

Planta annua, usque ad 40 cm alta, divaricato-ramosa, ascendens vel erecta. Caulis albo-villosus, pili patentes, c. 0.5-0.75 mm longi, plerumque ad bas in tuberculati. Folia lata elliptica vel ovata vel elliptico-ovata vel suborbicularia, 1.5-4.7 cm longa, 1-3 cm lata, apice obtusa, ad basin subtruncata, distince ad 2 m petiolata; pili supra appressi, vix tuberculati,

subtus patentes, semper tuberculati, nervis supra paulo impressis, subglabris, subtus paulo prominentibus, margine crassiuscule vel inconspicue revoluta. Inflorescentiae laterales vel terminales; cincinni singuli, unilaterales et uniseriati, laxiusculi, usque ad 17 cm longi, c. 10-25-flori. Calyces sessiles vel usque ad 2.5 mm pedicellati, laciniis linearo-lanceolatis, fere liberis, 2.5-4 mm longis, 0.5-1 mm latis, subacutis, extus dense subpatentivillosis, intus ad 2/3 superiora pilosis, 1/3 inferum subglabris. Corolla hypocrateriformis, 8-12 mm longa; tubus 5-7 mm, intus glaber, rarissimus pilosus, extus patenti-villosus; lobi explanati, rotundati, 1.5-2.5 mm lati, margine undulato-crenulati, in sinibus inter lobos semper dentati, aestivatione imbricati. Antherae (1.5) 1.8-2.2 mm longae, ad 1-1.5 mm supra corollae basin insertae, apice subito acutatae, paulo incurvatae. Stigma conico-elongatum, 1.2-1.7 mm longum, ad medium articulatam, apice bilobum, pilosum, in parte infera subglabrum. Stylus 0-0.5 mm, glaber. Nuculae ellipsoideae, 0.9-1.3 x 1.8-2 mm, glabrae, laeviusculae vel paulo lineolatae.

*H. esfandiarii* differt ab *H. szovitsii* (STEV.) BUNGE pilis patentibus (nec appressis), inflorescentii singulis (nec  $\pm$  ramosis), corolla hypocrateriformi (nec infundibuliformi), corollae lobis semper undulato-crenulatis (nec integris vel raro crenulatis); antheris longioribus (1.5) 1.8-2.2 mm longis (nec. 1-1.7 (1.8) longis).

Typus: Iran; Semnan Province: 15 km northeast of Garmsar, gravelly soils, 1000 m, 11.6.1990, H. AKHANI, 6359 (holo: MMIT, Natural History Museum of Iran; iso: W).

***Heliotropium Esfandiarii* AKHANI & H. RIEDL spec.nov. (Fig.1)**

Annual plant, up to 40 cm high, divaricately branched, ascending or erect. Stem whitish villose, hairs patent, c. 0.5-0.75 mm, at the base mostly tuberculate. Leaves elliptic or ovate or elliptic-ovate or suborbicular, 1.5-4.7 cm long and 1-3 cm broad, apex obtuse, at the base subtruncate, with a long petiole up to 2 cm; upper surface with subappressed hairs,  $\pm$  tuberculate at base; lower surface with dense tuberculate patent hairs; nerves in lower surface  $\pm$  prominent, in upper surface slightly impressed, subglabrous; margin slightly thickened, sometimes obscurely revolute. Inflorescence lateral and terminal; cymes single, up to 17 cm, rather lax with c. 10-25 unilateral, uniseriate, flowers. Calyx sessile or up to 2.5 mm pedicellate; calyx lobes linear-lanceolate, nearly free; 2.5-4 x 0.8-1 mm, subacute, outside with dense spreading hairs, inside in the upper 2/3 hairy, in the lower 1/3 subglabrous. Corolla hypocrateriform, 8-12 mm

long, tube 5-7 mm, inside glabrous, very rarely hairy, outside with patent hairs, lobes broad, rotundate, 1.5-2.5 mm broad, at the margin undulate-crenulate, with intercalary teeth. Anthers inserted 1-1.5 mm above corolla base, (1.5) 1.8-2.2 mm long; apex suddenly acute, rather curved inside. Stigma  $\pm$  elongate-conical, 1.2-1.7 mm, articulate at the middle, apex bilobed, pilose, lower part subglabrous. Style 0-0.5 mm, glabrous. Nutlets ellipsoid, 0.9-1.3 x 1.8-2 mm, glabrous, surface smooth to obscurely lineolate.

**D i a g n o s e :** *H. Esfandiarii* differs from *H. Szovitsii* (STEV.) BUNGE, by patent hairs (not appressed), single inflorescence (not  $\pm$  branched), hypocrateriform (not infundibular) corolla, always undulate-crenulate (not entire and sometimes crenulate) corolla lobes, (1.5) 1.8-2.2 mm  $\sqrt{\text{not } 1-1.7 \text{ (1.8)}}$  long anthers.

**M a t e r i a l s e e n :** Semnan: 15 km northeast of Garmsar, gravelly soils, 1000 m, 11.6.1990, H. AKHANI, 6359 (MMTT, type); road from Semnan to Firuzkuh, 15-25 km from Semnan, 1400 m, 27.9.1974, HEDGE, WENDELBO, ASSADI & ALA, 14731 (TARI); c. 10 km NW of Semnan, 26.7.1982, ASSADI & MOZAFFARIAN, 40359 (TARI); road of Garmsar towards Semnan, 11 km west of Deh-Namak, 800 m, 12.6.1990, H. AKHANI, 6366 (MMTT); 67 km from Semnan, 7.6.1972, AMIN & MOUSSAVI, 1278 (TARI); 8 km N Garmsar, Booneh-kuh, 19.6.1974, AMIN & BAZARGAN, 19050 (TARI); c. 15 km E of Garmsar, 1000 m, 27.9.1974, HEDGE, WENDELBO, ASSADI & ALA, 14765 (TARI); 50 km after Garmsar towards Semnan, 900 m, 7.7.1972, IRANSHAHR, 16265 (HMIA); Simindasht towards Kabutar-darreh, Hable-rud, 3.-4.9.1948, AELLEN & MANUCHEHRI, 5356 (W, HMIA); 13 km east of Eivanakey, gypsum and salted hills, 1000 m, 8.7.1974, RENZ & IRANSHAHR, 16575 (HMIA); along the river Hableh-rud, between Mahmoudabad and Kabutar-darreh, 1500 m, 5.9.1988, ASSADI & ABOUHAMZEH, 66337 (TARI); between Simindasht and Kabutardarreh, 1150 m, AELLEN, 1181 (W); Tehran: Varamin, Sharifabad, 1050 m, 12.10.1972, F. ROWSHAN, 6375 (TARI).

Among the examined plants there is one specimen (HEDGE, WENDELBO; ASSADI & ALA, 14765) the corolla of which is pilose inside. This shows that presence of hairs inside of corolla isn't a constant character in some species of the genus *Heliotropium*-contrary to earlier papers e.g. BUNGE 1869, POPOV 1953 & RIEDL 1967. According to recent investigations of AKHANI in some other species with their corolla usually glabrous inside,

there is a chance to find plants with pilose corolla inside. Such cases were found in *H. dissitiflorum*, *H. aucheri* and *H. dasycarpum*.

### Acknowledgments

We are much indebted to Mr. Moradi, Tehran for preparing the illustration. H. Akhane gratefully acknowledges the help of Mr. Ammarloei director of Natural History Museum of Iran (Department of Environment) during the investigation.

### References

- BUNGE, A., 1869: Über die *Heliotropium*-Arten der mittelländischen orientalischen Floren. - Bull. Soc. Imp. Naturalistes Moscou, 42(1), 279-332.
- POPOV, M.G., 1953: *Boraginaceae* in Flora URSS. 19: 97-691. Moscow, Leningrad.
- RIEDL, H., 1967: *Boraginaceae* in K.H. RECHINGER (ed), *Flora Iranica*, 48. Graz.

Address of the authors: Hossein A k h a n e

	Natural History Museum of Iran
	Department of Botany
	9-Ghaem-Magham Farahani Avenue
15899 TEHRAN	15899 TEHRAN
Iran	Iran

Dr. Harald R i e d l  
Vienna Natural History Museum  
Department of Botany  
Burgring 7  
A-1014 VIENNA  
Austria





Fig.1: *Heliotropium esfandiarii* AKHANI & H. RIEDL



Map 1: Distribution map of *Heliotropium esfandiarii* AKHANI & H. RIEDL